

Overwhelm vs Burnout

A Call Me Courage guide

This guide is for reflection and practical support, not diagnosis. Use it to notice patterns, name what may be happening, and consider a kind next step.

Why this distinction matters

Overwhelm can feel like too much arriving at once. Burnout is more often what develops when stress continues and your capacity to respond stays depleted. Telling them apart can help you choose support that fits the moment, without blame or shame.

How to use this guide

- **Notice** what is happening with curiosity rather than criticism.
- **Compare** the patterns, knowing that people can experience both.
- **Choose** a small, supportive next step instead of demanding a perfect solution.

Call me Courage

Your strength lies within!

The Courage Journey Method™
Overwhelm vs Burnout

Understanding Overwhelm

Overwhelm is the experience of having more demands, information, emotion, or stimulation than you can comfortably process at that moment. It can happen quickly, and it does not mean you are weak or incapable.

Common signs

Area	You might notice
Thoughts	Racing thoughts, forgetfulness, indecision, or difficulty focusing on one thing.
Emotions	Irritability, worry, tearfulness, frustration, or feeling unusually sensitive.
Body	Tension, a restless feeling, headaches, a fast heartbeat, shallow breathing, or sensory overload.
Behaviour	Procrastinating, rushing, withdrawing, snapping, multitasking, or finding it hard to start.

Common short-term triggers

- Several deadlines, requests, or decisions arriving together.
- A sudden change, uncertainty, conflict, or unexpected responsibility.
- Poor sleep, skipped meals, pain, illness, or a busy sensory environment.
- A build-up of smaller demands that leaves no space to reset.

Reflection prompt

What feels like the loudest demand right now, and what could be paused, shared, simplified, or left until later?

Understanding Burnout

Burnout is a pattern of prolonged stress in which your available capacity becomes depleted. It can affect how you think, feel, function, and connect with others. It is not simply ordinary tiredness, and it is not defined by one difficult day.

Key distinction

Overwhelm often says, “There is too much happening right now.” Burnout may feel more like, “I have been carrying too much for too long, and recovery is no longer catching up.” People can move between the two, and this description is not a medical diagnosis.

Possible signs over time

Area	You might notice
Energy	Persistent exhaustion, reduced reserves, or needing much longer to recover.
Motivation	Tasks that once felt manageable seem unusually hard to begin or complete.
Body	Sleep changes, muscle tension, digestive discomfort, headaches, or other stress-related sensations.

Work or study	Reduced concentration, more mistakes, avoidance, or feeling unable to meet usual demands.
Relationships	Withdrawing, having less patience, or feeling too depleted to stay connected.

Reflection Prompt

Which changes have lasted beyond a busy day, and what would it look like to treat that pattern as a signal for support rather than a personal failure?

Helping yourself

When overwhelm is high

1. Reduce the immediate load: silence non-essential alerts, step away from extra input, and write down what can wait.
2. Choose one next step that takes a few minutes. Make it smaller than your first instinct, such as opening the document, drinking water, or sending one honest message.
3. Give your body a reset: soften your shoulders, take a slower breath, change rooms, or sit somewhere quieter.
4. Name the need without arguing with it: “I am overloaded right now, so I am going to do less for the next ten minutes.”

When depletion has been lasting

☐ **Reduce demands** where possible. Renegotiate a deadline, remove a non-essential task, or ask what can be shared.
☐ **Protect recovery** with regular pauses, sleep-supportive routines, food, movement that feels manageable, and genuinely restorative time.
☐ **Set one boundary** around availability, workload, messages, or commitments. A clear “not today” can create breathing room.
☐ **Reconnect with support** by telling a trusted person what has been difficult and naming the kind of help that would be useful.
☐ **Seek qualified professional help** when symptoms persist, worsen, or significantly affect daily life, work, study, sleep, or relationships.

Helping someone else

You do not need to diagnose someone or solve everything. A calm, respectful presence can make it easier for them to notice what they need and choose support.

A supportive approach

1. Notice gently. Look for changes in energy, mood, participation, communication, or usual routines without making assumptions.
2. Approach with care. Choose a private moment and ask whether they have capacity to talk.
3. Listen before advising. Let pauses happen, reflect back what you hear, and avoid turning the conversation into an investigation.
4. Validate the experience. You can acknowledge that something is hard without agreeing with every conclusion or claiming to know exactly how they feel.
5. Offer specific practical help. Suggest one concrete option, such as taking a task, bringing food, helping draft a message, or sitting nearby while they begin.
6. Respect autonomy. Ask what would help, accept “no,” and avoid taking control of decisions that belong to them.
7. Follow up without taking over. A brief check-in can communicate care, while leaving room for their pace and privacy.

Supportive phrases

- “I have noticed things seem heavy lately. Would you like company or space?”

- “That sounds like a lot to carry. I am glad you told me.”
- “What would make today five percent easier?”
- “I can take care of this one task, if that would help.”

Phrases to avoid

- “Everyone feels like this.”
- “Just push through it.”
- “You need to be more organised.”
- “I know exactly how you feel.”

Urgent reminder

If someone may be in immediate danger, treat the situation as urgent and connect with appropriate emergency or urgent local support. You do not have to manage that concern alone.

Sources and further support

This guide is a concise, evidence-informed wellbeing resource built around noticing stress patterns, reducing demands, restoring recovery, strengthening connection, and seeking help when needed. For further support, consider trusted public-health, workplace, education, community, or qualified professional resources that fit your circumstances.

Recovery is not a test of willpower. You are always allowed to take a smaller step, change the course of your plan, and accept support before everything feels unmanageable. If you need help navigating overwhelm or burnout, please explore the resources listed below. In the footer of this document, you will find a link to the Call Me Courage Blog, which includes a dedicated crisis line and support listings for both the **USA** and **Canada**. If you are located outside these countries, please search your local

area for regional crisis lines or mental health support services to get the guidance you deserve.